

COSMOLOGICAL EVOLUTION OF THE DUTY CYCLE OF QUASARS

JIAN-MIN WANG¹, YAN-MEI CHEN^{1,2} AND FAN ZHANG¹
Received 2006 May 15; accepted 2006 June 28

ABSTRACT

Quasars are powered by accretion onto supermassive black holes, but the problem of the duty cycle related to the episodic activity of the black holes remains open as one of the major questions of cosmological evolution of quasars. In this Letter, we obtain quasar duty cycles based on analyses of a large sample composed of 10,979 quasars with redshifts $z \leq 2.1$ from the Sloan Digital Sky Survey (SDSS) Data Release Three. We estimate masses of quasar black holes and obtain their mass function (MF) of the present sample. We then get the duty cycle $\bar{\delta}(z) = 10^{-3} \sim 1$ based on the Soltan's argument, implying that black holes are undergoing multiple episodic activity. We find that the duty cycle has a strong evolution. By comparison, we show that evolution of the duty cycle follows the history of cosmic star formation rate (SFR) density in the Universe, providing intriguing evidence for a natural connection between star formation and triggering of black hole activity. Feedback on star formation from black hole activity is briefly discussed.

Subject headings: black hole physics — galaxies: active — galaxies: evolution — galaxies: nuclei — quasar: general

1. INTRODUCTION

Supermassive black holes are relics of quasars in the Universe (Soltan 1982; Rees 1984, 1990). Evolution of quasars is led by switching on and off accretion onto the black holes. During their entire evolution, how many times and how many black holes are triggered? what is the triggering mechanism? and why do quasars switch off?

The duty cycle, defined as the fraction of active black holes to their total number is a key to tackling the above problems (Richstone et al. 1998; Martini 2004). A popular method to get the duty cycle invokes the continuity equation and the MF of quasar black holes. With an assumption that quasar black holes have the same Eddington ratio, their MF can be obtained from the luminosity function and finally the duty cycle can be found from the continuity equation (Small & Blandford 1990; Marconi et al. 2004). This is a convenient way to discuss evolution of the black holes, but the degeneracy of the Eddington ratio and the duty cycle still holds. Actually, not only are the Eddington ratios *not* constant for different quasars at different epochs, but they appear to be quite scattered (Vestergaard 2004). The duty cycle is poorly understood as a statistical parameter tracing the evolution of quasar populations.

In recent years, there has been much progress in estimating black hole masses both in nearby galaxies and distant quasars. The empirical relation of reverberation mapping allows us to conveniently obtain the black hole masses from a large sample and directly get their MF. Thus it becomes realistic to get new clues to understand the evolution of quasars. Fortunately, by invoking the MF, we can decouple the degeneracy of the duty cycle and the Eddington ratio to get the duty cycle.

In this Letter, we use available SDSS data to directly get the MF of the black holes so as to discuss the duty cycle based on the MF and find that there is a strong cosmological evolution. Our calculations assume a cosmology with the Hubble constant $H_0 = 70 \text{ Mpc}^{-1} \text{ km s}^{-1}$, $\Omega_m = 0.3$, and $\Omega_\Lambda = 0.7$.

2. BLACK HOLE EVOLUTION

Active black hole evolution can be described by the MF $\Phi(M_\bullet, z)$, which is defined as $\Phi(M_\bullet, z) = d^2N/dM_\bullet dV$, where $M_\bullet$ is the black hole mass, dN is the number of quasars within the comoving volume element dV and mass interval $dM_\bullet$. The number density of quasar black holes is then given by $N_{\text{qso}}(z) = \int_{M_\bullet^*} \Phi(M_\bullet, z) dM_\bullet$ at redshift z , where $M_\bullet^*$ is the flux-limited black hole mass in a survey. If $\mathcal{N}(M_\bullet, z)$ is the MF of all black holes including the active and the inactive ones, the total number density of black holes is $N_{\text{tot}}(z) = \int_{M_\bullet^*} \mathcal{N}(M_\bullet, z) dM_\bullet$ at redshift z . The relation between $\mathcal{N}(M_\bullet, z)$ and $\Phi(M_\bullet, z)$ can be written as (Small & Blandford 1992; Marconi et al. 2004),

$$\Phi(M_\bullet, z) = \delta(M_\bullet, z) \mathcal{N}(M_\bullet, z). \quad (1)$$

where the duty cycle, $\delta(M_\bullet, z)$ is a function of black hole mass and redshift. Integrating equation (1) over all black hole masses, we have a mean duty cycle in terms of their number density

$$\bar{\delta}_1(z) = \frac{\int_{M_\bullet^*} \Phi(M_\bullet, z) dM_\bullet}{\int_{M_\bullet^*} \mathcal{N}(M_\bullet, z) dM_\bullet} = \frac{N_{\text{qso}}(z)}{N_{\text{tot}}(z)}. \quad (2)$$

This duty cycle represents the relative number of the active black holes to the total. Multiplying by $M_\bullet$ and integrating Eq. (1) over all black hole masses, we have an averaged duty cycle weighted by the masses of the black holes

$$\bar{\delta}_2(z) = \frac{\int_{M_\bullet^*} \Phi(M_\bullet, z) M_\bullet dM_\bullet}{\int_{M_\bullet^*} \mathcal{N}(M_\bullet, z) M_\bullet dM_\bullet} = \frac{\langle M_\bullet(z) \rangle_{\text{qso}}}{\langle M_\bullet(z) \rangle_{\text{all}}} \bar{\delta}_1(z), \quad (3)$$

where $\langle M_\bullet(z) \rangle_{\text{qso}} = \int_{M_\bullet^*} \Phi(M_\bullet, z) M_\bullet dM_\bullet / N_{\text{qso}}$ is the averaged mass of the quasar black holes at redshift z and $\langle M_\bullet(z) \rangle_{\text{all}} = \int_{M_\bullet^*} \mathcal{N}(M_\bullet, z) M_\bullet dM_\bullet / N_{\text{tot}}$ for all of active and inactive black holes. We show below $\langle M_\bullet(z) \rangle_{\text{qso}} = \langle M_\bullet(z) \rangle_{\text{all}}$ for $M_\bullet > M_\bullet^*$. So we have $\bar{\delta}_1(z) = \bar{\delta}_2(z)$.

The averaged mass of all black holes is $\langle M_\bullet(t) \rangle_{\text{all}} = \langle M_\bullet^S \rangle + \int_0^t \langle \dot{M}_\bullet(t') \rangle_{\text{all}} dt'$, where $\langle M_\bullet^S \rangle$ is the mean mass of seed black holes and $\langle \dot{M}_\bullet(t') \rangle_{\text{all}}$ is the averaged accretion rate of all black holes larger than $M_\bullet^*$. Considering that $\langle \dot{M}_\bullet(t') \rangle_{\text{all}} =$

¹ Key Laboratory for Particle Astrophysics, Institute of High Energy Physics, Chinese Academy of Sciences, 19B Yuquan Road, Beijing 100049, China

² Graduate School, Chinese Academy of Science, 19A Yuquan Road, Beijing 100049, China

$\bar{\delta}_1(t')\langle\dot{M}_\bullet(t')\rangle_{\text{qso}}$ [see Eq. (9) in Small & Blandford 1992; and the first equation in sentence with Eq. (14) in Marconi et al. 2004], we have $\langle M_\bullet(t) \rangle_{\text{all}} = \langle M_\bullet^S \rangle + \int_0^t \bar{\delta}_1(t')\langle\dot{M}_\bullet(t')\rangle_{\text{qso}} dt'$. On the other hand, the duty cycle can be written as $\bar{\delta}_1(t) = \Delta t_{\text{act}} / \Delta t$ for a single episodic phase within a time interval Δt , where $\Delta t = \Delta t_{\text{act}} + \Delta t_{\text{dor}}$; Δt_{act} and Δt_{dor} are the active and the dormant times, respectively. Then, for the i -th episodic phase within $\Delta t'_i$, the accreted mass is given by $\langle\dot{M}_\bullet(t'_i)\rangle_{\text{qso}} \bar{\delta}_1(t'_i) \Delta t'_i$. The mean mass of quasar black holes then reads $\langle M_\bullet(t) \rangle_{\text{qso}} = \langle M_\bullet^S \rangle + \sum_i \langle\dot{M}_\bullet(t'_i)\rangle_{\text{qso}} \bar{\delta}_1(t'_i) \Delta t'_i = \langle M_\bullet^S \rangle + \int_0^t \langle\dot{M}_\bullet(t')\rangle_{\text{qso}} \bar{\delta}_1(t') dt'$ after multiple episodic phases. We then have $\langle M_\bullet(t) \rangle_{\text{all}} = \langle M_\bullet(t) \rangle_{\text{qso}}$. This relation can be understood more easily if we consider multiple episodic growth of black holes. We thus have the duty cycle of the black holes in a concise form of $\bar{\delta}(z) \equiv \bar{\delta}_1(z) = \bar{\delta}_2(z)$.

The total mass density of the black holes at redshift z contributed from accretion is then given by

$$\rho_{\text{acc}}(z) = \int_{-\infty}^z \left(\frac{dt}{dz} \right) dz \int_{L_*(z)}^{\infty} \frac{1-\eta}{\eta} \frac{L_{\text{Bol}}}{c^2} \Psi(L, z) dL, \quad (4)$$

where $L_*(z)$ is the limit luminosity due to survey sensitivity, $\Psi(L, z)$ is the luminosity function, the bolometric luminosity is given by $L_{\text{Bol}} = C_B L_B$, with C_B the correction factor relating B -band luminosity L_B to L_{Bol} , η the radiative efficiency and c the light speed. This density includes *all* the black holes that were brighter than $L_*(z)$. The mass density of the active black holes at z with $L > L_*$ is given by

$$\rho_{\text{qso}}(z) = \int_{M_*^C}^{\infty} \Phi(M_\bullet, z) M_\bullet dM_\bullet, \quad (5)$$

where M_*^C is the limit black hole mass due to the limit luminosity. We apply the Sołtan's argument to the present sample at any redshift z

$$\rho_{\text{tot}}(z) = \int_{M_*^C}^{\infty} \mathcal{N}(M_\bullet, z) M_\bullet dM_\bullet = \rho_{\text{acc}}(z). \quad (6)$$

Though we do not know the distribution $\mathcal{N}(M_\bullet, z)$, we know the mass density of all black holes from Sołtan's argument. Finally, we have duty cycle $\bar{\delta}(z)$,

$$\bar{\delta}(z) = \frac{\rho_{\text{qso}}(z)}{\rho_{\text{acc}}(z)}. \quad (7)$$

If we know $\Psi(L, z)$ and $\Phi(M_\bullet, z)$, we can easily get the duty cycle of quasars at redshift z . We have to stress that Eq. (7) neither needs an initial condition nor the assumption of a constant Eddington ratio, something which must be specified for in the alternative approach solving the continuity equation (e.g. $\bar{\delta} = 1$ at $z = 3$ in Marconi et al. 2004).

3. SAMPLE AND BLACK HOLE MASS FUNCTION

3.1. Spectrum Analysis

The largest quasar sample, given by Richards et al. (2006) from the SDSS DR 3, consists of 15,343 quasars from $z = 0$ to 5, which is complete and homogeneous for apparent magnitude $i = 15 - 19$. We only use those quasars (11,954) with $z \leq 2.1$ in the present paper. We subtract the continuum and iron emission (based on the iron template derived from I Zw 1 spectra, which is kindly provided by R. J. McLure from private communication 2006). We then fit H β and Mg II lines. For those with $z \leq 0.7$, four components are used to model the spectra: broad and narrow H β plus narrow [O III] 4959Å and

[O III] 5007Å. For others, we use one broad and one narrow component to model the Mg II line. Some objects are removed from the sample for one of three reasons; they have: 1) too poor spectra to fit due to low signal-noise ratio; 2) only narrow lines ($< 2000 \text{ km s}^{-1}$) or 3) serious absorption at the Mg II line. Finally we have 10,979 quasars available for estimating black hole masses with $z \leq 2.1$. The removed quasars reduce the completeness of the sample given in Table 1. The last bin (z_{10}) is poor because the quality of quasar spectra is not good enough to measure width of Mg II. A future paper will give a detailed description of estimating black hole masses and related issues (Chen et al. in preparation).

3.2. Black Hole Mass Function

We apply the latest version of the empirical relation of reverberation mapping (Kaspi et al. 2005; Vestergaard & Peterson 2006) to calculate the black hole mass in each quasar. For low redshift quasars with $z \leq 0.7$, we use the full-width-half-maximum of H β whereas we use Mg II for those with $0.7 < z \leq 2.1$ (McLure & Dunlop 2004). The scatter of the re-calibrated relation for the black hole masses is less than 0.4 dex, which is much improved. Figure 1a shows the mass distribution of the present sample. We fit the mass distribution of the black holes via least square method in a form of three power laws

$$\mathcal{F}(m_\bullet) = f_0 m_\bullet^{\alpha_1} \left(1 + \frac{m_\bullet}{m_{\bullet 1}} \right)^{-\alpha_2} \left(1 + \frac{m_\bullet}{m_{\bullet 2}} \right)^{-\alpha_3}, \quad (8)$$

where $m_\bullet$ is mass of black holes in unit of solar mass. This expression has the limits $\mathcal{F}(m_\bullet) \propto m_\bullet^{\alpha_1}$ for $m_\bullet \ll m_{\bullet 1}$, $\mathcal{F}(m_\bullet) \propto m_\bullet^{\alpha_1 - \alpha_2}$ for $m_{\bullet 1} \ll m_\bullet \ll m_{\bullet 2}$ and $\mathcal{F}(m_\bullet) \propto m_\bullet^{\alpha_1 - \alpha_2 - \alpha_3}$ for $m_\bullet \gg m_{\bullet 2}$. We obtain $f_0 = (2.70 \pm 0.35) \times 10^{-25}$, $m_{\bullet 1} = (4.14 \pm 0.56) \times 10^7$, $m_{\bullet 2} = (2.50 \pm 0.22) \times 10^9$, $\alpha_1 = 3.56 \pm 0.28$, $\alpha_2 = 2.87 \pm 0.25$ and $\alpha_3 = 2.71 \pm 0.07$. A significant break appears at $M_\bullet = 2.5 \times 10^9 M_\odot$ and then a steeper mass spectrum $\mathcal{F}(m_\bullet) \propto m_\bullet^{-2.02}$ follows the break mass, which is consistent with the maximum mass from the SDSS DR1 (McLure & Dunlop 2004).

TABLE 1 THE MASS FUNCTION

z-bin	Φ_* (10^{-6})	β_1	β_2	$M_\bullet^*$ ($10^9 M_\odot$)	$\chi_j^2/d.o.f.$	C (%)
z_1	48.1	-0.60	1.55	0.01	4.40/4	98
z_2	4.74	0.32	2.48	0.16	7.90/9	95
z_3	5.94	0.79	2.58	0.18	8.77/10	93
z_4	0.79	0.18	2.89	0.58	9.89/10	94
z_5	0.44	0.48	3.36	1.25	8.25/10	95
z_6	0.27	0.15	3.21	1.81	8.16/9	96
z_7	0.29	0.42	2.82	1.53	8.78/8	97
z_8	0.33	0.68	2.86	1.55	8.58/8	96
z_9	0.28	0.50	2.54	1.29	6.45/7	89
z_{10}	0.15	0.63	2.30	1.26	6.71/7	67

The last column is the completeness at each redshift bin in our sample. The redshift bin z_j is defined by $z_j = (j - 1/2)\Delta z$, where $j = 1, \dots, 10$. The fittings are not included one point in z_1 , z_4 , z_6 and $z_8 - z_{10}$ -bins, which significantly deviate from the trend of double power-laws.

We get the MF by dividing our sample into 10 redshift bins with an interval $\Delta z = 0.21$ and then into 20 black hole mass bins in each redshift bin. Figure 1b shows the MF. We find that the function can be well fitted by double power laws in the following form

$$\Phi(M_\bullet, z) = \Phi_* \left[\left(\frac{M_\bullet^*}{M_\bullet} \right)^{\beta_1} + \left(\frac{M_\bullet}{M_\bullet^*} \right)^{\beta_2} \right]^{-1}, \quad (9)$$

where β_1 , β_2 , Φ_* and M_*^* are constants. The peak mass is given by $M_{\bullet}^{\text{peak}} = (\beta_1/\beta_2)^{1/(\beta_1+\beta_2)} M_*^*$. We get the four parameters from the least χ -square method via minimizing $\chi_j^2 = \sum_i [\Phi(M_{\bullet,i}, z) - \Phi_{ij}]^2 / \sigma_{ij}^2$, where Φ_{ij} is the MF derived from the sample in mass $M_{\bullet,i}$ -bin and redshift z_j -bin. Since the averaged error bar in the luminosity function is about $\Delta\Psi/\Psi \sim 0.1$ (Richards et al. 2006), we take $\sigma_{ij} = 0.1\Phi_{ij}$ for the MF. Table 1 lists all the parameters for each redshift bin.




FIG. 1.— *a* shows the mass distribution of the black holes in our sample. The red line is the least square fit. The black hole mass function $\Phi(M_{\bullet}, z)$ at each redshift bin is shown in *b*. The lines represent the best fits given in Tab 1.

The MF shows an increasing peak mass toward high redshifts. The mass break at the low mass side is not real. It is caused by the survey flux limit. We have to point out that the MF is *not* complete for all black holes, but it is complete for our flux-limited criterion.

4. RESULTS

The *B*-band luminosity is converted from the absolute magnitude $M_i(z=2)$ via a relation of $M_B = M_i(z=2) + 0.804$ (Richards et al. 2006). It has been found that the factor C_B is not a function of redshift (Steffen et al. 2006). A more elaborate treatment gives $C_B = 6-7$ for quasars (Marconi et al. 2004). We use $C_B = 6.5$ throughout the redshift range in this paper. The luminosity function is taken from Richards et al. (2006). We take $M_{\bullet}^C$ from the minimum mass of the black holes in each redshift bin to calculate their mass density, ρ_{qso} (Eq. 5). This corresponds to the survey limit and is consistent with the accretion density ρ_{acc} (Eq. 4). The radiative efficiency $\eta \geq 0.1$ is reached by Yu & Tremaine (2002), $\eta \geq 0.15$ by Elvis et al. (2002) and $\eta = 0.15$ by Gammie et al. (2004), based on the Sothan's argument, the X-ray background and numerical simulations, respectively. Numerical calculations indicate that black holes are rotating with their maximum spin all the time during their evo-

lution if accretion is included (Volonteri et al 2005), which is supported by studies of SDSS quasars (Wang et al. 2006); then $\eta \approx 0.3$ without significant evolution. We thus calculate the duty cycle for two different radiative efficiencies $\eta = 0.1$ and 0.3.




FIG. 2.— *a*, *b* show the duty cycle of quasars as a function of redshift in our sample and the history of the SFR density, respectively. The SFR density is taken from Pérez-González et al. (2005).

Results are shown in Figure 2. First, quasars have duty cycle of $\bar{\delta}(z) = 10^{-3} \sim 1$ as shown in Fig 2a, indicating that black holes are undergoing active and dormant phases, namely episodic activity. The value of $\bar{\delta}(z) \rightarrow 1$ at $z \sim 2$ agrees with the assumption of $\delta = 1$ at $z = 3$ in Marconi et al. (2004). Second, the duty cycle is rapidly evolving from $z \approx 2$ to the local Universe. We find $\bar{\delta}(z) \propto z^{\gamma}$, where $\gamma \sim 2.5$ until $z = 1.5$, above which it tends to flatten. Third, as shown in Fig. 2a and 2b, the duty cycle connects quite naturally with the history of the SFR density as a consequence of co-evolution of galaxies and black holes. Massive, gas-rich mergers account not only for most of the star formation at $z \approx 2-3$, but they are probably also responsible for triggering major episodes of black hole activity (Di Matteo et al. 2005). Figures 2a and 2b show that the duty cycle follows the star formation history, implying evidence for intrinsic star-formation-triggered black hole activity. The higher the SFR density, the higher the triggering frequencies of the black hole activity. Last, we define $\mathcal{R} = \Delta t_{\text{act}} / \Delta t_{\text{dor}}$, then have $\mathcal{R} = \bar{\delta}(z) / [1 - \bar{\delta}(z)]$. Episodic activity of the black holes can be described by this parameter. When $\bar{\delta}(z) \rightarrow 1$, black holes have $\mathcal{R} \gg 1$, namely, the dormant black holes are frequently triggered at $z \sim 2$ by star formation. At that time, quasars look like long-lived phenomena because of $\mathcal{R} \gg 1$.

5. DISCUSSION AND SUMMARY

The evolution of quasars is jointly controlled by the triggering mechanism and accretion. The duty cycle is a key parameter to unveil the evolution of quasars. The results of the present pa-

per show a very strong cosmological evolution of quasar's duty cycle. The triggering history represented by $\bar{\delta}(z)$ is quite similar to the evolution of cosmic SFR density. This indicates that star formation may be the direct mechanism to trigger the activity of the black holes.

The duty cycle can be roughly justified from the galaxy luminosity function. According to the luminosity function of galaxies at $1.8 \leq z \leq 2.0$ (Dahlen et al. 2005), the galaxy number density is $n_G \approx 826 \text{ Gpc}^{-3}$ for galaxies brighter than R -band magnitude $M_R = -24$. This corresponds to the number density of galaxies with black hole mass larger than $10^9 M_\odot$ converted from $\log(M_{\text{BH}}/M_\odot) = -0.5M_R - 3$ (McLure & Dunlop 2001). Number density of quasars brighter than $M_i = -28$ (corresponding to a black hole with mass $> 10^9 M_\odot$ if it is accreting at the Eddington limit) is $n_Q \approx 175 \text{ Gpc}^{-3}$ based on the quasar luminosity function (Richards et al. 2006). We thus estimate a duty cycle of ~ 0.18 , which is roughly consistent with the present results (see Figure 2a).

The SFR density rises with redshift out to $z = 1.5$ and appears to be roughly flat between $z \approx 1.5$ and $z \approx 3.0$. This tendency

could be explained by strong feedback from activity of black holes to their host galaxies (Silk 2005; Di Matteo et al. 2005; Croton et al. 2006). With a balance between star formation and feedback in $z \sim 1.5 - 3.0$, the violent star formation is then suppressed. However, the SFR density is going to decrease with time due to a shortage of gas and the BH duty cycle follows this trend. To further confirm this, future work will focus on the dependence of the duty cycle on the black hole masses. It could show the feedback-dependence on the BH growth itself. Additionally the total accretion time (net lifetime) of black holes will be then obtained.

The anonymous referee is greatly acknowledged for pointing out a mathematical error in the paper. The authors are grateful to M. Z. Kong, R. Wang and R. J. McLure for help in SDSS data analysis, and L. C. Ho, S. N. Zhang, J. X. Wang and X. Y. Xia for useful discussions. This research is supported by the Natural Science Foundation of China through NSFC-10325313, 10233030 and 10521001.

REFERENCES

- Croton, D. J., et al., 2006, MNRAS, 365, 11
Dahlen, T., et al., 2005, ApJ, 631, 126
Elvis, M., Risaliti, G., & Zamorani, G., 2002, ApJ, 565, L75
Gammie, C. F., Shapiro, S. L., & McKinney, J. C., 2004, ApJ, 602, 312
Di Matteo, T., Springel, V., & Hernquist, L., 2005, Nature, 433, 604
Kaspi, S., et al., 2005, ApJ, 629, 674
Marconi, A., Risaliti, G., Gilli, R., Hunt, L. K., Maiolino, R., & Salvati, M., 2004, MNRAS, 351, 169
Martini, P., 2004, in Coevolution of Black Holes and Galaxies, from the Carnegie Observatories Centennial Symposia. by Cambridge University Press, Ed. L. C. Ho, p. 169
Martini, P., & Schneider, D., 2003, ApJ, 597, L109
McLure, R., & Dunlop, J. S., 2001, MNRAS, 327, 199
McLure, R., & Dunlop, J. S., 2004, MNRAS, 352, 1390
Pérez-González, P. G., et al. 2005, ApJ, 630, 82
Rees, M. J., 1984, ARA&A, 22, 471
Rees, M. J., 1990, Science, 247, 817
Richards, G. T., et al., 2006, AJ, 131, 2766
Richstone, D., et al., 1998, Nature, 395A, 14
Silk, J., 2005, MNRAS, 364, 1337
Small, T.A., & Blandford, R. D., 1992, MNRAS, 259, 725
Soltan, A., 1982, MNRAS, 200, 115
Steffen, A. T., et al., 2006, AJ, 131, 2826
Vestergaard, M., 2004, ApJ, 601, 676
Vestergaard, M., & Peterson, B. M., 2006, ApJ, 641, 689
Volonteri, M., Madau, P., Quataert, E., Rees, M. J., 2005, ApJ, 620, 69
Wang, J.-M., Chen, Y.-M., Ho, L. C., & McLure, R. J., 2006, ApJ, 642, L111
Yu, Q., & Tremaine, S., 2002, MNRAS, 335, 965